

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 February 2001 (08.02.2001)

PCT

(10) International Publication Number
WO 01/09341 A2

(51) International Patent Classification⁷: C12N 15/53,
15/54, 1/21, C12Q 1/32, 1/48, C12N 9/04, 9/10

KLÖTI, Andreas, S. [CH/US]; 6 Forest Creek Drive,
Durham, NC 27713 (US).

(21) International Application Number: PCT/EP00/07033

(74) Common Representative: BAYER AKTIENGE-
SELLSCHAFT; D-51368 Leverkusen (DE).

(22) International Filing Date: 21 July 2000 (21.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
199 35 967.9 30 July 1999 (30.07.1999) DE
09/449,335 24 November 1999 (24.11.1999) US

(71) Applicants (for all designated States except US): BAYER
AKTIENGESELLSCHAFT [DE/DE]; D-51368 Lev-
erkusen (DE). PARADIGM GENETICS, INC. [US/US];
104 Alexander Drive, Research Triangle Park, NC
27709-4528 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): HAIN, Rüdiger
[DE/DE]; Talstr. 53a, D-40764 Langenfeld (DE). TIET-
JEN, Klaus-Günther [DE/DE]; Am Alten Broich 98,
D-40764 Langenfeld (DE). BUSCH, Marco [DE/DE];
Benninghausen 59 b, D-51399 Burscheid (DE). MARTIN,
William, F. [US/DE]; Rilkestr. 13, D-41469 Neuss (DE).

Published:

— Without international search report and to be republished
upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



WO 01/09341 A2

(54) Title: METHOD OF DETERMINING THE ACTIVITY OF 1-DEOXY-D-XYLULOSE-5-PHOSPHATE REDUCTOISO-
MERASE AND 1-DEOXY-D-XYLULOSE-5-PHOSPHATE SYNTHASE

(57) Abstract: The invention relates to DNA which encodes *Arabidopsis* 1-deoxy-xylulose-5-phosphate reductoisomerase and to a
method of identifying modulators of the 1-deoxy-D-xylulose-5-phosphate reductoisomerase and 1-deoxy-D-xylulose-5-phosphate
synthase activity.

Method of determining the activity of 1-deoxy-D-xylulose-5-phosphate reductoisomerase and 1-deoxy-D-xylulose-5-phosphate synthase

5 The invention relates to DNA which encodes for Arabidopsis 1-deoxy-D-xylulose-5-phosphate reductoisomerase, and to a method of identifying modulators of an enzyme with 1-deoxy-D-xylulose-5-phosphate reductoisomerase activity and of an enzyme with 1-deoxy-D-xylulose-5-phosphate synthase activity.

10 Undesired vegetation can be prevented by using herbicides. The demands made of herbicides have risen constantly with regard to activity, costs and ecofriendliness. There is therefore a need for new substances which can be developed into new potent herbicides. In general, it is normal to search for such new guide structures in greenhouse tests. However, such tests are laborious and expensive. The number of substances which can be tested in the greenhouse is, accordingly, limited.

15 The search is underway for plant-specific biosynthetic routes which constitute advantageous sites of action for herbicides and do not occur in animal organisms.

20 Recently, it has been found that plants synthesize essential plastid isoprenoids not by exploiting the mevalonate biosynthetic pathway which is found in the animal organism, but the microbial 1-deoxy-xylulose-5-phosphate biosynthetic pathway (Lichtenthaler, K. (1998), *Fett/Lipid* 100, 128-138; Eisenreich et al). (1998), *Chemistry & Biology* 5, R221-R233).

25 This biosynthetic pathway finally leads to the synthesis of, *inter alia*, carotenoids: and the side chains of plastoquinone and chlorophyll. These products are essential for the photosynthetic growth of plants, Inhibition of one step in this biosynthetic pathway entails the end of plant growth.

30 This is why the 1-deoxy-xylulose-5-phosphate biosynthetic pathway is of particular interest in the search for new herbicidally active compounds. In particular the two

enzymes 1-deoxy-D-xylulose-5-phosphate synthase (DXPS) and 1-deoxy-D-xylulose-5-phosphate reductoisomerase (DXPR) are of central importance. It has already been demonstrated that DXPS (CLA I) is essential for the development of a normal plant (Mandel et al. (1996), Plant J. 9, 649-658). This discovery supports the expectation that a herbicidal compound which affects DXPS activity has a herbicidal action. Also, it has been demonstrated that bacterial DXFR is inhibited by the herbicidal compound Fosmidomycin, which is already known (Zeidler et al. (1998), Z. Naturforsch. 53, 980-986; Kuzuyama et al. (1998), Tetrahedron Lett. 39, 7913-7916). However, there are no commercially useable herbicides, which affect DXPS or DXPR activity. In the search for new, improved herbicides, both enzymes are therefore of high importance as sites of action. The 1-deoxy-xylulose-5-phosphate biosynthetic pathway also has importance in microorganisms, especially in parasitic microorganisms as, for example, bacteria or plasmodia. The treatment of infectious diseases, in particular the treatment of malaria, may be based on the inhibition of this metabolic pathway (Jomaa et al. (1999), Science 285, 1573-1576).

So far it has not been possible to determine the enzymatic activity of the enzyme 1-deoxy-xylulose-5-phosphate synthase in a simple test system since the reaction neither entails a measurable change in absorption nor can be coupled in a simple manner with a colour change or fluorescence change (Figure 1). The enzymatic activity of 1-deoxy-D-xylulose-5-phosphate reductoisomerase entails a measurable change in the optical absorption of the cosubstrate NADPH. However the substrate 1-deoxy-D-xylulose-5-phosphate can only be synthesized with difficulty (Figure 1). While 1-deoxy-D-xylulose-5-phosphate can be prepared via a chemical or biochemical route, both methods are expensive and are not well suited for use in test systems with high throughput (Taylor et al. (1998), J. Org. Chem. 63, 2375-2377; Blagg and Poulter (1999), J. Org. Chem. 64, 1508-1511).

The present invention solves these problems by combining the two enzymes in one test system. The *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate synthase gene is already known under the name CLAI (Mandel et al. (1996), Plant J. 5.

649-658). The 1-deoxy-D-xylulose-5-phosphate reductoisomerase gene is, as yet, known only from *Mentha piperita* (Lange et al. (1998), Proc. Natl. Acad. Sci. U.S.A. 5, 2100-2104) and from various microorganisms. The *A. thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase has also been described and a fragment of
5 the amino acid sequence has been published (Lange und Croteau (1999). Archives of Biochem. and Biophys. 365, 170-174). The combined reaction of pyruvate and glyceraldehyde-3-phosphate to give 2-C-methyl-D-erythrol-4-phosphate is monitored by visually detecting the NADPH consumption (Figure 1). The test system is suitable for the search for modulators of both enzymes, that is to say substances which inhibit
10 or else stimulate the activity of the enzymes, and can be used for test series with high throughput (high-throughput screening, HTS). After detection of modulators by one of the two enzymes in an HTS system, the modulators of the two enzymes can be distinguished from each other by using the existing methods for measuring the activity of the two enzymes (Sprenger et al. (1997), Proc. Natl. Acad. Sci. U.S.A. 94,
15 12857-12862; Kuzuyama et al. (1998), Tetrahedron Lett. 39, 4509-4512; DE 197 52 700-A1).

The present invention relates to the DNA which encodes *Arabidopsis* 1-deoxy-D-xylulose-5-phosphate reductoisomerase, in particular the *Arabidopsis thaliana*
20 1-deoxy-D-xylulose-5-phosphate reductoisomerase, and to fragments of this DNA, which encode functional subunits of 1-deoxy-D-xylulose-5-phosphate reductoisomerase.

The invention furthermore relates to DNA which encodes *Arabidopsis* 1-deoxy-D-xylulose-5-phosphate reductoisomerase, with an amino acid sequence as shown in
25 SEQ ID NO 2 or SEQ ID NO 6.

The invention furthermore relates to DNA as described under SEQ ID NO 1 or SEQ ID NO 5 which encodes the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate
30 reductoisomerase.

The invention furthermore relates to DNA which shows 80%, preferably 90%, homology to the DNA described under SEQ ID NO 1 or SEQ ID NO 5 and which encodes plant 1-deoxy-D-xylulose-5-phosphate reductoisomerase.

5 The invention furthermore relates to DNA which is complementary to the DNA which encodes the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase, and to RNA which is complementary to the DNA which encodes the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase.

10 The invention furthermore relates to an expression construct which encompasses DNA which encodes the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase and is described under SEQ ID NO 1 or SEQ ID NO 5, and to a sequence which is functionally linked herewith and which allows the 1-deoxy-D-xylulose-5-phosphate reductoisomerase to be expressed.

15 The invention furthermore relates to a vector which comprises DNA which encodes the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase and/or is described under SEQ ID NO 1 or SEQ ID NO 5 and which allows the 1-deoxy-D-xylulose-5-phosphate reductoisomerase to be expressed in a host cell.

20 The invention furthermore relates to a host cell which comprises the abovementioned DNA, an expression construct as mentioned above, or a vector which allows the 1-deoxy-D-xylulose-5-phosphate reductoisomerase to be expressed.

25 The invention furthermore relates to the use of DXPR and/or DXPS modulators as herbicides, antibiotic agents or anti-malarial agents.

The invention also relates to the use of DXPR and/or DXPS modulators as lead structures for the chemical optimization and the development of improved modulators.

30

The invention also relates to a method of determining the activity of 1-deoxy-D-xylulose-5-phosphate reductoisomerase and 1-deoxy-D-xylulose-5-phosphate synthase, which is based on combining the conversion of pyruvate and glyceraldehyde-3-phosphate to give 1-deoxy-D-xylulose-5-phosphate by 1-deoxy-D-xylulose-5-phosphate synthase with the conversion of the resulting 1-deoxy-D-xylulose-5-phosphate to give 2-C-methyl-D-erythrol-4-phosphate by 1-deoxy-D-xylulose-5-phosphate reductoisomerase in one test system.

The course of the overall reaction, that is to say the conversion of the pyruvate and glyceraldehyde-3-phosphate into 2-C-methyl-D-erythrol-4-phosphate is monitored with reference to the optical change by the decrease in the cofactor NADPH of 1-deoxy-D-xylulose-5-phosphate reductoisomerase.

The invention also relates to a method of identifying of substances which modify the activity of 1-deoxy-D-xylulose-5-phosphate reductoisomerase or 1-deoxy-D-xylulose-5-phosphate synthase, in which the above-described test system for determining the activity of 1-deoxy-D-xylulose-5-phosphate reductoisomerase and of 1-deoxy-D-xylulose-5-phosphate synthase is used. The test system is optimized in such a way that an optimal conversion of pyruvate and glyceraldehyde-3-phosphate to give 2-C-methyl-D-erythrol-4-phosphate is ensured. The reaction can be carried out in the presence and absence of substances which modify the activity of one of the enzymes involved. A comparison of the reaction in the presence and absence of such a substance with reference to the NADPH consumption thus allows substances which modulate, preferably inhibit, the activity of 1-deoxy-D-xylulose-5-phosphate reductoisomerase and/or of 1-deoxy-D-xylulose-5-phosphate synthase to be identified.

The invention also relates to substances which are found with the aid of the above described method, with the exception of Fosmidomycin, which is already known to inhibit 1-deoxy-D-xylulose-5-phosphate reductoisomerase (Zeidler et al. (1998), Z. Naturforsch. 53, 980-986).

The invention also relates to the use of substances which are found with the aid of the above-described method for use as modulators, preferably as inhibitors, of the enzymatic activity of 1-deoxy-D-xylulose-5-phosphate reductoisomerase and/or of 1-deoxy-D-xylulose-5-phosphate synthase.

5

The invention also relates to the use of substances which are found with the aid of the above-described method for use as herbicides, antibiotic agents or anti-malarial agents.

10

The term "functional fragments" describes those DNA fragments which encode polypeptides which still have 1-deoxy-D-xylulose-5-phosphate reductoisomerase activity, or fragments of 1-deoxy-D-xylulose-5-phosphate reductoisomerase which still have this activity.

15

The term "homology" in relation to DNA means that DNA segments which are at least 15 base pairs long or strands which are complementary to the DNA match the corresponding DNA in at least 80%, preferably in 90%, of the nucleotides. Such a homology is determined, inter alia, with the aid of computer programs such as the GCG program (Devereux et al. (1983), Nucleic Acids Res. 12, 3 87-395).

20

"Homology" exists also when a DNA segment is capable of hybridizing with the DNA strand in question or with its complementary strand.

25

The term "to hybridize" or "hybridization" describes the process in which a single stranded nucleic acid molecule undergoes base pairing with a complementary DNA strand, where the capability of a single-stranded nucleic acid molecule depends on the stringency of the hybridization conditions.

30

The term "stringency" relates to the hybridization conditions. "High stringency makes base pairing difficult. To do this, high temperatures of 42°C or less are used, a formamide concentration of less than 20% and low salt (SSC) concentrations, Alter-

natively, temperatures of 65°C or less can be used in combination with a low salt concentration (SSPE). "Low stringency" conditions favour the formation of base pairs. The temperatures used here are 37°C or less, the formamide concentration is less than 50%, and the salt concentration (SSC) is moderate. Alternatively, temperatures of 50°C or less in combination with a medium to high salt concentration (SSPE) are used.

The term "complementary" relates to the capability of purine and pyrimidine nucleotides to form base pairs with each other via hydrogen bonds. Complementary base pairs are, *inter alia*, guanine and cytosine, adenine and thymine, and adenine and uracil.

The term "plasmid" refers to an extrachromosomal genetic element. The original plasmids used for the present invention are either commercially available or freely accessible or can be derived from such plasmids by known methods.

The term "vector" describes a DNA element used for introducing exogenous DNA into host cells. A vector contains a nucleotide sequence which encodes one or more polypeptides.

One skilled in the art is aware of the fact that the degenerate genetic code (i.e. 64 codons encode 20 amino acids) allow a large number of "silent" substitutions of nucleotide base pairs to be introduced into the sequence shown here without changing the identity of the protein products encoded by it. The scope of the invention includes all such substitutions.

DNA isolation

The nucleic acid mentioned here can exist in complete cells, in cell lysates, in partially purified or biologically pure form, i.e. when other cell components or chemical

precursors and by-products, in the case of chemical DNA synthesis, have been removed.

5 The DNA mentioned here can be obtained by a series of genetic and recombinant DNA techniques, for example by means of amplification with the aid of the polymerase chain reaction (PCR) or else by de novo DNA synthesis. The DNA mentioned here can be isolated by means of PCR amplification of genomic DNA from suitable plant cells using oligonucleotide primers which are directed at a suitable region of SEQ ID NO 1 or SEQ ID NO 5 (see, for example, J. Sambrook et al, (1989), Molecular Cloning, 2nd edition, chapter 14).

Obtaining and purifying the protein

15 The invention also relates to polypeptides which have 1-deoxy-D-xylulose-5-phosphate reductoisomerase activity and which are encoded by an above-described DNA.

20 The skilled worker knows that the polypeptides of the present invention can be obtained by various routes, for example by chemical methods such as the solid-phase method. To obtain larger quantities of protein, the use of recombinant methods is recommended. Expression of a cloned 1-deoxy-D-xylulose-5-phosphate reductoisomerase gene or fragments thereof can take place in a series of suitable host cells which are known to the skilled worker. To this end, a 1-deoxy-D-xylulose-5-phosphate reductoisomerase gene is introduced into a host cell with the aid of known methods.

25 The integration of the cloned 1-deoxy-D-xylulose-5-phosphate reductoisomerase gene in the chromosome of the host cell is within the scope of the present invention. Preferably, the gene or fragments thereof are inserted into a plasmid, and the encoding regions of the 1-deoxy-D-xylulose-5-phosphate reductoisomerase gene or fragments thereof are functionally linked to a constitutive or inducible promoter.

The basic steps for generating the recombinant 1-deoxy-D-xylulose-5-phosphate reductoisomerase are:

- 5 1. Obtaining a natural, synthetic or semi-synthetic DNA which encodes
 1-deoxy-D-xylulose-5-phosphate reductoisomerase.
2. Introducing this DNA into an expression vector which is suitable for ex-
 pressing 1-deoxy-D-xylulose-5-phosphate reductoisomerase either alone or as
10 a fused protein.
3. Transformation of a suitable host cell, preferably a prokaryotic host cell, with
 this expression vector.
- 15 4. Growing this transformed host cell in a manner which is suitable for express-
 ing 1-deoxy-D-xylulose-5-phosphate reductoisomerase.
5. Harvesting the cells and purifying 1-deoxy-D-xylulose-5-phosphate reductoi-
 somerase by suitable known methods.

20

The encoding regions of 1-deoxy-D-xylulose-5-phosphate reductoisomerase and of 1-deoxy-D-xylulose-5-phosphate synthase can be expressed by the customary methods in *E. coli*, either separately or together. Suitable expression systems for *E. coli* are commercially available, for example the expression vectors of the pET series, for
25 example pET3a, pET23a, pET28a with his-Tag or pET32a with his-Tag for the simple purification and thioredoxin fusion for improving the solubility of the expressed enzyme, and pGEX with glutathion synthetase fusion. The expression vectors are transformed into XDE3-lysogenic *E. coli* strains, for example, BL21(DE3), HMS
174(DE3) or AD494(DE3). After the cells have become attached, expression is in-
30 duced with IPTG under standard conditions known to the skilled worker. After cell induction, incubation is carried out for 3 to 24 hours at temperatures from 18°C to

37°C. The cells are disrupted by sonication in disruption buffer (10 to 200 mM sodium phosphate, 100 to 500 mM NaCl, pH 5 to 8). The protein which has been expressed can be purified by chromatographic methods, in the case of protein which has been expressed with a his-Tag by means of chromatography on an Ni-NTA column.

Since, according to current knowledge, the 1-deoxy-xylulose-5-phosphate pathway does not exist in animals and yeasts, expression of the protein in commercially available yeast strains (for example *Pichia pastoris*) or in insect cell cultures (for example Sf9 cells) is another favourable option.

Alternatively, the proteins may also be expressed in plants.

Determination of the 1-deoxy-D-xylulose-5-phosphate reductoisomerase and/or 1-deoxy-D-xylulose-5-phosphate synthase activity, and identification of modulators of the enzyme activity of both enzymes

To identify and develop herbicidally active substances, it is necessary to find a way of determining the effect of various candidate substances on 1-deoxy-D-xylulose-5-phosphate reductoisomerase and 1-deoxy-D-xylulose-5-phosphate synthase activity. To this end, a way must be found of measuring the activity of both enzymes in an efficient and simple manner, or of detecting an inhibition or else stimulation of this activity.

One way of determining the effect of a substance on the enzymatic reaction of 1-deoxy-D-xylulose-5-phosphate reductoisomerase and/or of 1-deoxy-D-xylulose-5-phosphate synthase is to contact purified 1-deoxy-D-xylulose-5-phosphate reductoisomerase and 1-deoxy-D-xylulose-5-phosphate synthase, or fragments with 1-deoxy-D-xylulose-5-phosphate reductoisomerase activity and fragments with 1-deoxy-D-xylulose-5-phosphate synthase activity, with a test substance and to check the activity of both enzymes.

According to the present invention, the 1-deoxy-D-xylulose-5-phosphate reductoisomerase and 1-deoxy-D-xylulose-5-phosphate synthase activity is determined in a combined test system which contains both 1-deoxy-D-xylulose-5-phosphate reductoisomerase and 1-deoxy-D-xylulose-5-phosphate synthase. In this test system, 1-deoxy-D-xylulose-5-phosphate synthase converts pyruvate and glyceraldehyde-3-phosphate into 1-deoxy-D-xylulose-5-phosphate, which is converted by 1-deoxy-D-xylulose-5-phosphate reductoisomerase directly into 2-C-methyl-D-erythrol-4-phosphate, with consumption of NADPH. The decrease in the NADPH concentration can be monitored with aid of optical measurement methods (Figure 1).

To test for substances which modulate, preferably inhibit, the 1-deoxy-D-xylulose-5-phosphate reductoisomerase and/or 1-deoxy-D-xylulose-5-phosphate synthase activity, the test system and the enzyme concentrations are designed in such a way that an optimal conversion of the pyruvate and glyceraldehyde-3-phosphate to give 2-C-methyl-D-Erythrol-4-phosphate is ensured. If one of the enzymes involved is inhibited or activated by a candidate substance, this can be detected by a drop or increase in the NADPH conversion.

Then, a separate activity test for 1-deoxy-D-xylulose-5-phosphate reductoisomerase or 1-deoxy-D-xylulose-5-phosphate synthase may be carried out in the known manner in order to determine which of the two enzymes is affected in its activity by the substance which has been found.

As an alternative to the known, separate activity tests, HPLC makes it possible to determine, starting from the combined activity test, which products or intermediates starting from the materials pyruvate and glyceraldehyde-3-phosphate have been formed. If 1-deoxy-D-xylulose-5-phosphate has been formed, but not 2-C-methyl-D-erythrol-4-phosphate, then it was DXPR which was inhibited. If 1-deoxy-D-xylulose-5-phosphate has not been formed either, the DXPS was (also) inhibited.

Example 1

Construction of a vector for expressing the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate synthase

5

The encoding sequence of the *Arabidopsis thaliana* CLA1 gene from position +1 to +2154 was amplified with the aid of the PCR technique using the following primers of the sequences shown in SEQ ID NO. 3 and SEQ ID NO. 4 (Mandel et al. (1996) CLA1, a novel gene required for chloroplast development, is highly conserved in evolution; Plant J. 5 649-658; Lange M. et al. (1998) A family of transketolases that directs isoprenoid biosynthesis via a mevalonate-independent pathway; Proc. Natl. Acad. Sci., U. S. A. 95, 2100-2104).

The template used was *Arabidopsis thaliana* single-stranded cDNA from 4-week-old seedlings.

The amplified fragment which carries the encoding sequence of the *Arabidopsis thaliana* DXPS was then cleaved with the restriction enzymes *Bam*HI and *Not*I. The resulting *Bam*HI/*Not*I-DXPS fragment was ligated into the linearized and dephosphorylated bacterial expression vector pET32 a (+) (Novagen). The resulting construct pET32-DXPS contains the encoding DXPS sequence within the reading frame with a fragment of the bacterial thioredoxin gene.

Example 2

25

Construction of a vector for expressing the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase

The encoding sequence of the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase was amplified with the aid of the PCR technique using the primers with the sequences as shown in SEQ ID NO. 3 and SEQ ID NO. 4. The amplified

30

fragment was then cleaved with the restriction endonucleases *EcoRI* and *Sall*. The resulting *EcoRI/Sall*-DXR fragment was ligated into the linearized and dephosphorylated bacterial expression vector pET32 a (+) (Novagen). In the resulting construct pET32-DXR the encoding DXR sequence was within the reading frame together with a fragment of the bacterial thioredoxin gene.

Example 3

Test system for identifying 1-deoxy-D-xylulose-5-phosphate reductoisomerase and/or 1-deoxy-D-xylulose-5-phosphate synthase modulators in an HTS

A microtitre plate (96-well format) is filled with solutions of 1-deoxy-xylulose-5-phosphate synthase (0.1 - 10 µg of purified enzyme/100 µl) and 1-deoxy-xylulose-5-phosphate reductoisomerase (0.1 - 10 µg of purified enzyme/100 µl) and of the co-substrate NADPH (0.1 - 10 mM) in the customary buffer (10 - 200 mM sodium phosphate pH 5 - 8), containing thiamine diphosphate (0.1 - 10mM) and MgCl₂ (0.5 - 50 mM). All concentrations (also those given further below) are based on the concentration after addition of all assay components. The candidate chemical, or as control, buffer, is pipetted into each cavity of the microtitre plate. After addition of the substrates pyruvate (1 - 100 mM) and glyceraldehyde-3-phosphate (1 - 100 mM) in customary buffer (as above, but without NADPH and without thiamine diphosphate), the plates are incubated between 18°C and 45°C until a drop in the optical density of NADPH at 340 nm which can be measured easily has been reached. The optical density is then read in a customary microtitre plate reader. Substances which inhibit one of the two enzymes which participate are identified by a reduced drop of the NADPH concentration with the aid of the optical density measurement.

In microtitre plates of greater density (384-well, 1536-well format and the like), the volumes indicated above are adapted to suit the system.

Explanation of the figure and of the sequence data

Fig. I shows the conversion of pyruvate and glyceraldehyde-3-phosphate via 1-deoxy-D-xylulose-5-phosphate to give 2-C-methyl-D-erythrol-4-phosphate, which is catalysed by the enzymes 1-deoxy-D-xylulose-5-phosphate synthase and 1-deoxy-D-xylulose-5-phosphate reductoisomerase. The reaction which is catalysed by 1-deoxy-D-xylulose-5-phosphate reductoisomerase requires NADPH as cofactor. The catalysed reactions are essential for the isoprenoid synthesis in plants.

SEQ ID NO. 1

DNA sequence encoding the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase. The other sequence shown is the amino acid sequence encoded by the DNA.

SEQ ID NO. 2

Amino acid sequence of the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase with 477 amino acids.

SEQ ID NO. 3

Oligonucleotide derived from the encoding sequence of the *Arabidopsis thaliana* CLAI gene for amplifying the *Arabidopsis thaliana* CLAI gene by means of the PCR technique, including nucleotides for a BamH1 cloning site.

SEQ ID NO. 4

Oligonucleotide derived from the encoding sequence of the *Arabidopsis thaliana* CLAI gene for amplifying the *Arabidopsis thaliana* CLAI gene by means of the PCR technique, including nucleotides for a NotI cloning site.

SEQ ID NO. 5

Another DNA sequence encoding the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase. Change at nucleotides 890-892. The other sequence shown is the amino acid sequence encoded by the DNA.

5

SEQ ID NO. 6

Another amino acid sequence of the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase with 477 amino acids. Change at amino acid 292.

10

SEQ ID NO. 7

Oligonucleotide derived from the encoding sequence of the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase gene for amplifying the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase gene by means of the PCR technique.

15

SEQ ID NO. 8

Oligonucleotide derived from the encoding sequence of the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase gene for amplifying the *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase gene by means of the PCR technique.

20


Patent Claims

1. DNA which encodes *Arabidopsis* 1-deoxy-D-xylulose-5-phosphate reductoisomerase or functional fragments thereof.
- 5 2. DNA according to Claim 1, characterized in that the DNA encodes the total, or functional fragments of an amino acid sequence as shown in SEQ ID NO. 2 or SEQ ID NO. 6.
- 10 3. DNA according to Claim 1 or 2, characterized in that the DNA encodes *Arabidopsis thaliana* 1-deoxy-D-xylulose-5-phosphate reductoisomerase.
4. DNA according to any of Claims 1 to 3 with a sequence as shown in SEQ ID NO. 1, SEQ ID NO. 5 or fragments thereof.
- 15 5. DNA, characterized in that its homology to the sequence as shown in SEQ ID NO. 1 or SEQ ID NO. 5 is at least 80%.
- 20 6. DNA according to Claim 5, characterized in that its homology to the sequence as shown in SEQ ID NO. 1 or SEQ ID NO. 5 is at least 90%.
7. DNA, characterized in that it is complementary to the DNA according to any of Claims 1 to 6.
- 25 8. RNA, characterized in that it is complementary to the DNA according to any of Claims 1 to 6.
- 30 9. Expression construct, characterized in that it encompasses DNA as claimed in any of Claims 1 to 6 and a sequence which is functionally linked herewith and which allows the DNA to be expressed.

10. Vector, characterized in that it contains DNA according to any of Claims 1 to 7 and 9.
- 5 11. Host cell containing DNA according to any of Claims 1 to 7, an expression construct according to Claim 9 or a vector according to Claim 10.
12. Use of 1-deoxy-D-xylulose-5-phosphate reductoisomerase and/or 1-deoxy-D-xylulose-5-phosphate synthase modulators as herbicides, antibiotic agents or anti-malarial agents.
- 10 13. Method of determining the activity of 1-deoxy-D-xylulose-5-phosphate reductoisomerase and/or 1-deoxy-D-xylulose-5-phosphate synthase, characterized in that
- 15 (a) pyruvate and glyceraldehyde-3-phosphate are converted into 1-deoxy-D-xylulose-5-phosphate with the aid of 1-deoxy-D-xylulose-5-phosphate synthase, and
- (b) the resulting 1-deoxy-D-xylulose-5-phosphate is converted into
- 20 2-C-methyl-D-erythrol-4-phosphate with the aid of 1-deoxy-D-xylulose-5-phosphate reductoisomerase.
14. Method according to Claim 13, characterized in that the conversion of 1-deoxy-D-xylulose-5-phosphate into 2-C-methyl-D-erythrol-4-phosphate is
- 25 measured with reference of the NADPH consumed during this conversion.
15. Method according to Claim 13 or 14, characterized in that the 1-deoxy-D-xylulose-5-phosphate reductoisomerase activity is determined.
- 30 16. Method according to Claim 13 or 14, characterized in that the 1-deoxy-D-xylulose-5-phosphate synthase activity is determined.

17. Method of identifying substances which modify the activity of 1-deoxy-D-xylulose-5-phosphate reductoisomerase and/or 1-deoxy-D-xylulose-5-phosphate synthase, characterized in that the method according to any of Claims 13 to 16 is carried out in the presence and absence of a test substance, and the activities found in each case are compared with each other.
18. Substances which are identified by a method according to Claim 17, with the exception of Fosmidomycin.
19. Use of substances which are identified by a method according to Claim 17 as modulators of the enzymatic activity of 1-deoxy-D-xylulose-5-phosphate reductoisomerase and/or of 1-deoxy-D-xylulose-5-phosphate synthase.
20. The use of substances identified by a method according to Claim 18 as herbicides, antibiotic agents or anti-malarial agents.
21. Polypeptide represented by SEQ ID NO. 2 or SEQ ID NO. 6.

1/1

**Figure 1**

SEQUENCE LISTING

<110> Busch, Marco
Hain, Rudiger
Martin, William
Tietjen, Klaus
Kloti, Andreas

<120> Method of determining the activity of
1-deoxy-D-xylulose-5-phosphate reductoisomerase and
1-deoxy-D-xylulose-5-phosphate synthase

<130> 2020US

<140>

<141>

<150> DE 199 35 967.9

<151> 1999-07-30

<160> 8

<170> PatentIn Ver. 2.1

<210> 1

<211> 1732

<212> DNA

<213> Arabidopsis thaliana

<220>

<221> CDS

<222> (18)..(1448)

<223> DNA encoding 1-deoxy-D-xylulose-5-phosphate
reductoisomerase

<400> 1

```

ttttttaaaa gactctg atg atg aca tta aac tca cta tct cca gct gaa   50
                Met Met Thr Leu Asn Ser Leu Ser Pro Ala Glu
                  1             5             10

tcc aaa gct att tct ttc ttg gat acc tcc agg ttc aat cca atc cct   98
Ser Lys Ala Ile Ser Phe Leu Asp Thr Ser Arg Phe Asn Pro Ile Pro
              15             20             25

aaa ctc tca ggt ggg ttt agt ttg agg agg agg aat caa ggg aga ggt   146
Lys Leu Ser Gly Gly Phe Ser Leu Arg Arg Arg Asn Gln Gly Arg Gly
              30             35             40

ttt gga aaa ggt gtt aag tgt tca gtg aaa gtg cag cag caa caa caa   194
Phe Gly Lys Gly Val Lys Cys Ser Val Lys Val Gln Gln Gln Gln
              45             50             55

cct cct cca gca tgg cct ggg aga gct gtc cct gag gcg cct cgt caa   242
Pro Pro Pro Ala Trp Pro Gly Arg Ala Val Pro Glu Ala Pro Arg Gln
              60             65             70             75

tct tgg gat gga cca aaa ccc atc tct atc gtt gga tct act ggt tct   290
Ser Trp Asp Gly Pro Lys Pro Ile Ser Ile Val Gly Ser Thr Gly Ser
              80             85             90

att ggc act cag aca ttg gat att gtg gct gag aat cct gac aaa ttc   338
Ile Gly Thr Gln Thr Leu Asp Ile Val Ala Glu Asn Pro Asp Lys Phe
              95             100             105

```

aga gtt gtg gct cta gct gct ggt tcg aat gtt act cta ctt gct gat	386
Arg Val Val Ala Leu Ala Ala Gly Ser Asn Val Thr Leu Leu Ala Asp	
110 115 120	
cag gta agg aga ttt aag cct gca ttg gtt gct gtt aga aac gag tca	434
Gln Val Arg Arg Phe Lys Pro Ala Leu Val Ala Val Arg Asn Glu Ser	
125 130 135	
ctg att aat gag ctt aaa gag gct tta gct gat ttg gac tat aaa ctc	482
Leu Ile Asn Glu Leu Lys Glu Ala Leu Ala Asp Leu Asp Tyr Lys Leu	
140 145 150 155	
gag att att cca gga gag caa gga gtg att gag gtt gcc cga cat cct	530
Glu Ile Ile Pro Glu Glu Gln Gly Val Ile Glu Val Ala Arg His Pro	
160 165 170	
gaa gct gta acc gtt gtt acc gga ata gta ggt tgt gcg gga cta aag	578
Glu Ala Val Thr Val Val Thr Gly Ile Val Gly Cys Ala Gly Leu Lys	
175 180 185	
cct acg gtt gct gca att gaa gca gga aag gac att gct ctt gca aac	626
Pro Thr Val Ala Ala Ile Glu Ala Gly Lys Asp Ile Ala Leu Ala Asn	
190 195 200	
aaa gag aca tta atc gca ggt ggt cct ttc gtg ctt ccg ctt gcc aac	674
Lys Glu Thr Leu Ile Ala Gly Gly Pro Phe Val Leu Pro Leu Ala Asn	
205 210 215	
aaa cat aat gta aag att ctt ccg gca gat tca gaa cat tct gcc ata	722
Lys His Asn Val Lys Ile Leu Pro Ala Asp Ser Glu His Ser Ala Ile	
220 225 230 235	
ttt cag tgt att caa ggt ttg cct gaa ggc gct ctg cgc aag ata atc	770
Phe Gln Cys Ile Gln Gly Leu Pro Glu Gly Ala Leu Arg Lys Ile Ile	
240 245 250	
ttg act gca tct ggt gga gct ttt agg gat tgg cct gtc gaa aag cta	818
Leu Thr Ala Ser Gly Gly Ala Phe Arg Asp Trp Pro Val Glu Lys Leu	
255 260 265	
aag gaa gtt aaa gta gcg gat gcg ttg aag cat cca aac tgg aac atg	866
Lys Glu Val Lys Val Ala Asp Ala Leu Lys His Pro Asn Trp Asn Met	
270 275 280	
gga aag aaa atc act gtg gac tcc tgt acg ctt ttc aac aag ggt ctt	914
Gly Lys Lys Ile Thr Val Ser Cys Thr Leu Phe Asn Lys Gly Leu	
285 290 295	
gag gtc att gaa gcg cat tat ttg ttt gga gct gag tat gac gat ata	962
Glu Val Ile Glu Ala His Tyr Leu Phe Gly Ala Glu Tyr Asp Asp Ile	
300 305 310 315	
gag att gtc att cat ccg caa agt atc ata cat tcc atg att gaa aca	1010
Glu Ile Val Ile His Pro Gln Ser Ile Ile His Ser Met Ile Glu Thr	
320 325 330	
cag gat tca tct gtg ctt gct caa ttg ggt tgg cct gat atg cgt tta	1058
Gln Asp Ser Ser Val Leu Ala Gln Leu Gly Trp Pro Asp Met Arg Leu	
335 340 345	

```

ccg att ctc tac acc atg tca tgg ccc gat aga gtt cct tgt tct gaa 1106
Pro Ile Leu Tyr Thr Met Ser Trp Pro Asp Arg Val Pro Cys Ser Glu
      350                      355                      360

gta act tgg cca aga ctt gac ctt tgc aaa ctc ggt tca ttg act ttc 1154
Val Thr Trp Pro Arg Leu Asp Leu Cys Lys Leu Gly Ser Leu Thr Phe
      365                      370                      375

aag aaa cca gac aat gtg aaa tac cca tcc atg gat ctt gct tat gct 1202
Lys Lys Pro Asp Asn Val Lys Tyr Pro Ser Met Asp Leu Ala Tyr Ala
      380                      385                      390

gct gga cga gct gga ggc aca atg act gga gtt ctc agc gcc gcc aat 1250
Ala Gly Arg Ala Gly Gly Thr Met Thr Gly Val Leu Ser Ala Ala Asn
      400                      405                      410

gag aaa gct gtt gaa atg ttc att gat gaa aag ata agc tat ttg gat 1298
Glu Lys Ala Val Glu Met Phe Ile Asp Glu Lys Ile Ser Tyr Leu Asp
      415                      420                      425

atc ttc aag gtt gtg gaa tta aca tgc gat aaa cat cga aac gag ttg 1346
Ile Phe Lys Val Val Glu Leu Thr Cys Asp Lys His Arg Asn Glu Leu
      430                      435                      440

gta aca tca ccg tct ctt gaa gag att gtt cac tat gac ttg tgg gca 1394
Val Thr Ser Pro Ser Leu Glu Glu Ile Val His Tyr Asp Leu Trp Ala
      445                      450                      455

cgt gaa tat gcc gcg aat gtg cag ctt tct tct ggt gct agg cca gtt 1442
Arg Glu Tyr Ala Ala Asn Val Gln Leu Ser Ser Gly Ala Arg Pro Val
      460                      465                      470                      475

cat gca tgaagaattg gttgttgga gaacataagg aagcttctga ggaaatgttg 1498
His Ala

aaagaagatt agtgtagaga atgggggtact acttaatagc gtttttggca aggattatgg 1558

attgtgtagc taatttatct gtgatccgaa caagccaaac tgataatttg aaaccattgt 1618

gatccgaaca agccaaactg ataatttgaa accattttta ccaataaaaac cgagcttaat 1678

tgtttcacat tatatgatta attacattca tctaagggtt cttgaaaaaa aaaa 1732

```

<210> 2

<211> 477

<212> PRT

<213> Arabidopsis thaliana

<400> 2

```

Met Met Thr Leu Asn Ser Leu Ser Pro Ala Glu Ser Lys Ala Ile Ser
  1                      5                      10                      15

Phe Leu Asp Thr Ser Arg Phe Asn Pro Ile Pro Lys Leu Ser Gly Gly
      20                      25                      30

Phe Ser Leu Arg Arg Arg Asn Gln Gly Arg Gly Phe Gly Lys Gly Val
      35                      40                      45

Lys Cys Ser Val Lys Val Gln Gln Gln Gln Gln Pro Pro Pro Ala Trp
      50                      55                      60

```

Pro Gly Arg Ala Val Pro Glu Ala Pro Arg Gln Ser Trp Asp Gly Pro
 65 70 75 80
 Lys Pro Ile Ser Ile Val Gly Ser Thr Gly Ser Ile Gly Thr Gln Thr
 85 90 95
 Leu Asp Ile Val Ala Glu Asn Pro Asp Lys Phe Arg Val Val Ala Leu
 100 105 110
 Ala Ala Gly Ser Asn Val Thr Leu Leu Ala Asp Gln Val Arg Arg Phe
 115 120 125
 Lys Pro Ala Leu Val Ala Val Arg Asn Glu Ser Leu Ile Asn Glu Leu
 130 135 140
 Lys Glu Ala Leu Ala Asp Leu Asp Tyr Lys Leu Glu Ile Ile Pro Gly
 145 150 155 160
 Glu Gln Gly Val Ile Glu Val Ala Arg His Pro Glu Ala Val Thr Val
 165 170 175
 Val Thr Gly Ile Val Gly Cys Ala Gly Leu Lys Pro Thr Val Ala Ala
 180 185 190
 Ile Glu Ala Gly Lys Asp Ile Ala Leu Ala Asn Lys Glu Thr Leu Ile
 195 200 205
 Ala Gly Gly Pro Phe Val Leu Pro Leu Ala Asn Lys His Asn Val Lys
 210 215 220
 Ile Leu Pro Ala Asp Ser Glu His Ser Ala Ile Phe Gln Cys Ile Gln
 225 230 235 240
 Gly Leu Pro Glu Gly Ala Leu Arg Lys Ile Ile Leu Thr Ala Ser Gly
 245 250 255
 Gly Ala Phe Arg Asp Trp Pro Val Glu Lys Leu Lys Glu Val Lys Val
 260 265 270
 Ala Asp Ala Leu Lys His Pro Asn Trp Asn Met Gly Lys Lys Ile Thr
 275 280 285
 Val Asp Ser Cys Thr Leu Phe Asn Lys Gly Leu Glu Val Ile Glu Ala
 290 295 300
 His Tyr Leu Phe Gly Ala Glu Tyr Asp Asp Ile Glu Ile Val Ile His
 305 310 315 320
 Pro Gln Ser Ile Ile His Ser Met Ile Glu Thr Gln Asp Ser Ser Val
 325 330 335
 Leu Ala Gln Leu Gly Trp Pro Asp Met Arg Leu Pro Ile Leu Tyr Thr
 340 345 350
 Met Ser Trp Pro Asp Arg Val Pro Cys Ser Glu Val Thr Trp Pro Arg
 355 360 365
 Leu Asp Leu Cys Lys Leu Gly Ser Leu Thr Phe Lys Lys Pro Asp Asn
 370 375 380
 Val Lys Tyr Pro Ser Met Asp Leu Ala Tyr Ala Ala Gly Arg Ala Gly
 385 390 395 400

Asn Val Gln Leu Ser Ser Gly Ala Arg Pro Val His Ala
465 470 475

t t t g g a a a g g t g t t a a g t g t c a g t g a a a g t g c a g c a g c a a c a a c a a 194
Phe Gly Lys Gly Val Lys Cys Ser Val Lys Val Gln Gln Gln Gln Gln
45 50 55

cct cct cca gca tgg cct ggg aga gct gtc cct gag gcg cct cgt caa	242
Pro Pro Pro Ala Trp Pro Gly Arg Ala Val Pro Glu Ala Pro Arg Gln	
60 65 70 75	
tct tgg gat gga cca aaa ccc atc tct atc gtt gga tct act ggt tct	290
Ser Trp Asp Gly Pro Lys Pro Ile Ser Ile Val Gly Ser Thr Gly Ser	
80 85 90	
att ggc act cag aca ttg gat att gtg gct gag aat cct gac aaa ttc	338
Ile Gly Thr Gln Thr Leu Asp Ile Val Ala Glu Asn Pro Asp Lys Phe	
95 100 105	
aga gtt gtg gct cta gct gct ggt tgc aat gtt act cta ctt gct gat	386
Arg Val Val Ala Leu Ala Ala Gly Ser Asn Val Thr Leu Leu Ala Asp	
110 115 120	
cag gta agg aga ttt aag cct gca ttg gtt gct gtt aga aac gag tca	434
Gln Val Arg Arg Phe Lys Pro Ala Leu Val Ala Val Arg Asn Glu Ser	
125 130 135	
ctg att aat gag ctt aaa gag gct tta gct gat ttg gac tat aaa ctc	482
Leu Ile Asn Glu Leu Lys Glu Ala Leu Ala Asp Leu Asp Tyr Lys Leu	
140 145 150 155	
gag att att cca gga gag caa gga gtg att gag gtt gcc cga cat cct	530
Glu Ile Ile Pro Gly Glu Gln Gly Val Ile Glu Val Ala Arg His Pro	
160 165 170	
gaa gct gta acc gtt gtt acc gga ata gta ggt tgt gcg gga cta aag	578
Glu Ala Val Thr Val Val Thr Gly Ile Val Gly Cys Ala Gly Leu Lys	
175 180 185	
cct acg gtt gct gca att gaa gca gga aag gac att gct ctt gca aac	626
Pro Thr Val Ala Ala Ile Glu Ala Gly Lys Asp Ile Ala Leu Ala Asn	
190 195 200	
aaa gag aca tta atc gca ggt ggt cct ttc gtg ctt ccg ctt gcc aac	674
Lys Glu Thr Leu Ile Ala Gly Gly Pro Phe Val Leu Pro Leu Ala Asn	
205 210 215	
aaa cat aat gta aag att ctt ccg gca gat tca gaa cat tct gcc ata	722
Lys His Asn Val Lys Ile Leu Pro Ala Asp Ser Glu His Ser Ala Ile	
220 225 230 235	
ttt cag tgt att caa ggt ttg cct gaa ggc gct ctg cgc aag ata atc	770
Phe Gln Cys Ile Gln Gly Leu Pro Glu Gly Ala Leu Arg Lys Ile Ile	
240 245 250	
ttg act gca tct ggt gga gct ttt agg gat tgg cct gtc gaa aag cta	818
Leu Thr Ala Ser Gly Gly Ala Phe Arg Asp Trp Pro Val Glu Lys Leu	
255 260 265	
aag gaa gtt aaa gta gcg gat gcg ttg aag cat cca aac tgg aac atg	866
Lys Glu Val Lys Val Ala Asp Ala Leu Lys His Pro Asn Trp Asn Met	
270 275 280	
gga aag aaa atc act gtg gac tct gct acg ctt ttc aac aag ggt ctt	914
Gly Lys Lys Ile Thr Val Asp Ser Ala Thr Leu Phe Asn Lys Gly Leu	
285 290 295	
gag gtc att gaa gcg cat tat ttg ttt gga gct gag tat gac gat ata	962
Glu Val Ile Glu Ala His Tyr Leu Phe Gly Ala Glu Tyr Asp Asp Ile	
300 305 310 315	

gag att gtc att cat ccg caa agt atc ata cat tcc atg att gaa aca 1010
 Glu Ile Val Ile His Pro Gln Ser Ile Ile His Ser Met Ile Glu Thr
 320 325 330

cag gat tca tct gtg ctt gct caa ttg ggt tgg cct gat atg cgt tta 1058
 Gln Asp Ser Ser Val Leu Ala Gln Leu Gly Trp Pro Asp Met Arg Leu
 335 340 345

ccg att ctc tac acc atg tca tgg ccc gat aga gtt cct tgt tct gaa 1106
 Pro Ile Leu Tyr Thr Met Ser Trp Pro Asp Arg Val Pro Cys Ser Glu
 350 355 360

gta act tgg cca aga ctt gac ctt tgc aaa ctc ggt tca ttg act ttc 1154
 Val Thr Trp Pro Arg Leu Asp Leu Cys Lys Leu Gly Ser Leu Thr Phe
 365 370 375

aag aaa cca gac aat gtg aaa tac cca tcc atg gat ctt gct tat gct 1202
 Lys Lys Pro Asp Asn Val Lys Tyr Pro Ser Met Asp Leu Ala Tyr Ala
 380 385 390 395

gct gga cga gct gga ggc aca atg act gga gtt ctc agc gcc gcc aat 1250
 Ala Gly Arg Ala Gly Gly Thr Met Thr Gly Val Leu Ser Ala Ala Asn
 400 405 410

gag aaa gct gtt gaa atg ttc att gat gaa aag ata agc tat ttg gat 1298
 Glu Lys Ala Val Glu Met Phe Ile Asp Glu Lys Ile Ser Tyr Leu Asp
 415 420 425

atc ttc aag gtt gtg gaa tta aca tgc gat aaa cat cga aac gag ttg 1346
 Ile Phe Lys Val Val Glu Leu Thr Cys Asp Lys His Arg Asn Glu Leu
 430 435 440

gta aca tca ccg tct ctt gaa gag att gtt cac tat gac ttg tgg gca 1394
 Val Thr Ser Pro Ser Leu Glu Glu Ile Val His Tyr Asp Leu Trp Ala
 445 450 455

cgt gaa tat gcc gcg aat gtg cag ctt tct tct ggt gct agg cca gtt 1442
 Arg Glu Tyr Ala Ala Asn Val Gln Leu Ser Ser Gly Ala Arg Pro Val
 460 465 470 475

cat gca tgaagaattg gttgttgga gaacataagg aagcttctga ggaaatgttg 1498
 His Ala

aaagaagatt agttagaga atgggtact acttaatagc gtttttggca aggattatgg 1558

attgtgtagc taatttatct gtgatccgaa caagccaaac tgataatttg aaaccattgt 1618

gatccgaaca agccaaactg ataatttgaa accattttta ccaataaaaac cgagcttaat 1678

tgtttcacat tatatgatta attacattca tctaagggtt cttgaaaaaa aaaa 1732

<210> 6

<211> 477

<212> PRT

<213> Arabidopsis thaliana

<400> 6

Met Met Thr Leu Asn Ser Leu Ser Pro Ala Glu Ser Lys Ala Ile Ser

1

5

10

15

Phe Leu Asp Thr Ser Arg Phe Asn Pro Ile Pro Lys Leu Ser Gly Gly
 20 25 30
 Phe Ser Leu Arg Arg Arg Asn Gln Gly Arg Gly Phe Gly Lys Gly Val
 35 40 45
 Lys Cys Ser Val Lys Val Gln Gln Gln Gln Gln Pro Pro Pro Ala Trp
 50 55 60
 Pro Gly Arg Ala Val Pro Glu Ala Pro Arg Gln Ser Trp Asp Gly Pro
 65 70 75 80
 Lys Pro Ile Ser Ile Val Gly Ser Thr Gly Ser Ile Gly Thr Gln Thr
 85 90 95
 Leu Asp Ile Val Ala Glu Asn Pro Asp Lys Phe Arg Val Val Ala Leu
 100 105 110
 Ala Ala Gly Ser Asn Val Thr Leu Leu Ala Asp Gln Val Arg Arg Phe
 115 120 125
 Lys Pro Ala Leu Val Ala Val Arg Asn Glu Ser Leu Ile Asn Glu Leu
 130 135 140
 Lys Glu Ala Leu Ala Asp Leu Asp Tyr Lys Leu Glu Ile Ile Pro Gly
 145 150 155 160
 Glu Gln Gly Val Ile Glu Val Ala Arg His Pro Glu Ala Val Thr Val
 165 170 175
 Val Thr Gly Ile Val Gly Cys Ala Gly Leu Lys Pro Thr Val Ala Ala
 180 185 190
 Ile Glu Ala Gly Lys Asp Ile Ala Leu Ala Asn Lys Glu Thr Leu Ile
 195 200 205
 Ala Gly Gly Pro Phe Val Leu Pro Leu Ala Asn Lys His Asn Val Lys
 210 215 220
 Ile Leu Pro Ala Asp Ser Glu His Ser Ala Ile Phe Gln Cys Ile Gln
 225 230 235 240
 Gly Leu Pro Glu Gly Ala Leu Arg Lys Ile Ile Leu Thr Ala Ser Gly
 245 250 255
 Gly Ala Phe Arg Asp Trp Pro Val Glu Lys Leu Lys Glu Val Lys Val
 260 265 270
 Ala Asp Ala Leu Lys His Pro Asn Trp Asn Met Gly Lys Lys Ile Thr
 275 280 285
 Val Asp Ser Ala Thr Leu Phe Asn Lys Gly Leu Glu Val Ile Glu Ala
 290 295 300
 His Tyr Leu Phe Gly Ala Glu Tyr Asp Asp Ile Glu Ile Val Ile His
 305 310 315 320
 Pro Gln Ser Ile Ile His Ser Met Ile Glu Thr Gln Asp Ser Ser Val
 325 330 335
 Leu Ala Gln Leu Gly Trp Pro Asp Met Arg Leu Pro Ile Leu Tyr Thr
 340 345 350

Met Ser Trp Pro Asp Arg Val Pro Cys Ser Glu Val Thr Trp Pro Arg
 355 360 365
 Leu Asp Leu Cys Lys Leu Gly Ser Leu Thr Phe Lys Lys Pro Asp Asn
 370 375 380
 Val Lys Tyr Pro Ser Met Asp Leu Ala Tyr Ala Ala Gly Arg Ala Gly
 385 390 395 400
 Gly Thr Met Thr Gly Val Leu Ser Ala Ala Asn Glu Lys Ala Val Glu
 405 410 415
 Met Phe Ile Asp Glu Lys Ile Ser Tyr Leu Asp Ile Phe Lys Val Val
 420 425 430
 Glu Leu Thr Cys Asp Lys His Arg Asn Glu Leu Val Thr Ser Pro Ser
 435 440 445
 Leu Glu Glu Ile Val His Tyr Asp Leu Trp Ala Arg Glu Tyr Ala Ala
 450 455 460
 Asn Val Gln Leu Ser Ser Gly Ala Arg Pro Val His Ala
 465 470 475

<210> 7

<211> 32

<212> DNA

<213> Arabidopsis thaliana

<400> 7

cctagaattc atgacattaa actcactatc tc

32

<210> 8

<211> 30

<212> DNA

<213> Arabidopsis thaliana

<400> 8

atatgtcgac tcatgcatga actggcctag

30

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 February 2001 (08.02.2001)

PCT

(10) International Publication Number
WO 01/09341 A3

(51) International Patent Classification⁷: C12N 15/53,
15/54, 1/21, C12Q 1/32, 1/48, C12N 9/04, 9/10

KLÖTI, Andreas, S. [CH/US]; 6 Forest Creek Drive,
Durham, NC 27713 (US).

(21) International Application Number: PCT/EP00/07033

(74) Common Representative: BAYER AKTIENGE-
SELLSCHAFT; D-51368 Leverkusen (DE).

(22) International Filing Date: 21 July 2000 (21.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
199 35 967.9 30 July 1999 (30.07.1999) DE
09/449,335 24 November 1999 (24.11.1999) US

(71) Applicants (for all designated States except US): BAYER
AKTIENGESELLSCHAFT [DE/DE]; D-51368 Lev-
erkusen (DE). PARADIGM GENETICS, INC. [US/US];
104 Alexander Drive, Research Triangle Park, NC
27709-4528 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(72) Inventors; and

(75) Inventors/Applicants (for US only): HAIN, Rüdiger
[DE/DE]; Talstr. 53a, D-40764 Langenfeld (DE). TIET-
JEN, Klaus-Günther [DE/DE]; Am Alten Broich 98,
D-40764 Langenfeld (DE). BUSCH, Marco [DE/DE];
Benninghausen 59 b, D-51399 Burscheid (DE). MARTIN,
William, F. [US/DE]; Rilkestr. 13, D-41469 Neuss (DE).

(88) Date of publication of the international search report:
23 August 2001

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



WO 01/09341 A3

(54) Title: METHOD OF DETERMINING THE ACTIVITY OF 1-DEOXY-D-XYLULOSE-5-PHOSPHATE REDUCTOISO-
MERASE AND 1-DEOXY-D-XYLULOSE-5-PHOSPHATE SYNTHASE

(57) Abstract: The invention relates to DNA which encodes *Arabidopsis* 1-deoxy-xylulose-5-phosphate reductoisomerase and to a
method of identifying modulators of the 1-deoxy-D-xylulose-5-phosphate reductoisomerase and 1-deoxy-D-xylulose-5-phosphate
synthase activity.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 00/07033

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/53 C12N15/54 C12N1/21 C12Q1/32 C12Q1/48
C12N9/04 C12N9/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE, STRAND, CAB Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMPLN 'Online! EMBL Heidelberg, Germany; AC AJ242588, 21 May 1999 (1999-05-21) SCHWENDER J: "Cloning and heterologous expression of a cDNA encoding 1-deoxy-d-xylulose-5-phosphate reductoisomerase of Arabidopsis thaliana" XP002146168 abstract	1-7
X	-& SCHWENDER J ET AL.: "Cloning and heterologous expression of a cDNA encoding 1-deoxy-D-xylulose-5-phosphate reductoisomerase of Arabidopsis thaliana" FEBS LETTERS, vol. 455, 16 July 1999 (1999-07-16), pages 140-144, XP002162455 the whole document --- -/-	8-11,21



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

9 March 2001

Date of mailing of the international search report

21/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Oderwald, H

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 00/07033

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>KUZUYAMA T ET AL: "Fosmidomycin, a Specific Inhibitor of 1-Deoxy-d-Xylulose 5-Phosphate Reductoisomerase in the Nonmevalonate Pathway for Terpenoid Biosynthesis"</p> <p>TETRAHEDRON LETTERS,NL,ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM,</p> <p>vol. 39, no. 43,</p> <p>22 October 1998 (1998-10-22), pages 7913-7916, XP004137840</p> <p>ISSN: 0040-4039</p> <p>cited in the application</p> <p>the whole document</p> <p style="text-align: center;">---</p>	<p>12-15,</p> <p>17,19,20</p>
A	<p>DE 298 00 547 U (HOECHST SCHERING AGREVO GMBH ;KERNFORSCHUNGSANLAGE JUELICH (DE))</p> <p>8 April 1999 (1999-04-08)</p> <p>the whole document</p> <p style="text-align: center;">---</p>	<p>13,14,</p> <p>16,17</p>
P,X	<p>WO 00 42205 A (NOVARTIS ERFINDE VERWALT GMBH ;NOVARTIS AG (CH); BUDZISZEWSKI GREGO) 20 July 2000 (2000-07-20)</p> <p>abstract; claims 1-46; examples 5,6E,7,8E,9</p> <p>page 12 -page 23</p> <p>page 27 -page 48</p> <p style="text-align: center;">---</p>	<p>1-12,</p> <p>17-21</p>
P,X	<p>WO 00 34448 A (DU PONT ;LEE JIAN MING (US); TAO YONG (US); CAHOON REBECCA E (US)) 15 June 2000 (2000-06-15)</p> <p>the whole document</p> <p style="text-align: center;">---</p>	<p>1-11,21</p>
E	<p>WO 00 63389 A (BORONAT ALBERT ;BHAT B GANESH (US); CALGENE LLC (US); KISHORE GANE) 26 October 2000 (2000-10-26)</p> <p>the whole document</p> <p style="text-align: center;">-----</p>	<p>1-11,21</p>

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-11, 21

DNA which encodes Arabidopsis 1-deoxy-D-xylulose-5-phosphate reductoisomerase or functional fragments thereof. RNA complementary to said DNA, expression construct, vector, host containing said DNA. Polypeptide represented by SEQ ID NO: 2 or 6.

2. Claims: 12, 17-20

Use of 1-deoxy-D-xylulose-5-phosphate reductoisomerase and/or 1-deoxy-D-xylulose-5-phosphate synthase modulators as herbicides, antibiotic agents or anti-malarial agents. Method of identifying substances which modify the activity of 1-deoxy-D-xylulose-5-phosphate reductoisomerase and/or 1-deoxy-D-xylulose-5-phosphate synthase, substances identified by said method, uses of said substances.

3. Claims: 13-16

Method of determining the activity of 1-deoxy-D-xylulose-5-phosphate reductoisomerase and/or 1-deoxy-D-xylulose-5-phosphate synthase.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 12, 18-20 (partially)

Present claims 12, 18-20 relate to a product/compound defined by reference to a desirable characteristic or property, namely modulator of DXR and/or DXS.

The claims cover all products/compounds having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such products/compounds. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the product/compound by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the product/compound fosmidomycin.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/07033

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 29800547 U	08-04-1999	DE 19752700 A JP 11169186 A	02-06-1999 29-06-1999
WO 0042205 A	20-07-2000	AU 2290900 A	01-08-2000
WO 0034448 A	15-06-2000	AU 2163300 A	26-06-2000
WO 0063389 A	26-10-2000	AU 4249300 A AU 4357900 A WO 0063391 A	02-11-2000 02-11-2000 26-10-2000